

### Datasheet for ABIN7646227

# anti-SLIT1 antibody



#### Overview

Quantity:	100 μL
Target:	SLIT1
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SLIT1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)
Product Details	
Purpose:	Polyclonal Antibody to Slit Homolog 1 (Slit1)
Immunogen:	RPD354Ra03Recombinant Slit Homolog 1 (Slit1)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against Slit1. It has been selected for its ability to recognize Slit1 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	
Target:	SLIT1
Alternative Name:	Slit1 (SLIT1 Products)

#### **Target Details**

- Angel Detaile		
Background:	MEGF4, SLIL1, SLIT3, Multiple epidermal growth factor-like domains protein 4	
UniProt:	088279	
Pathways:	Regulation of Cell Size	
Application Details		
Application Notes:	Western blotting: 0.5-5 µg/mL,Immunohistochemistry: 5-50 µg/mL,Immunocytochemistry: 5-	

Application Notes.	western blotting. 0.5-5 μg/mz,immunonistochemistry. 5-50 μg/mz,immunocytochemistry. 5-
	50 μg/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	500 μg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.