

Datasheet for ABIN7646357
anti-SLC19A1 antibody



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	SLC19A1
Reactivity:	Rat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This SLC19A1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP)

Product Details

Purpose:	Monoclonal Antibody to Folate Transporter (FOLT)
Immunogen:	CPE435Ra210VA Conjugated Folate Transporter (FOLT)
Clone:	C1
Specificity:	The antibody is a mouse monoclonal antibody raised against FOLT. It has been selected for its ability to recognize FOLT in immunohistochemical staining and western blotting.
Cross-Reactivity:	Rat
Purification:	Protein A + Protein G affinity chromatography

Target Details

Target:	SLC19A1
---------	---------

Target Details

Alternative Name:	FOLT (SLC19A1 Products)
Background:	SLC19A1, RFC1, CHMD, IFC1, REFC, Solute Carrier Family 19 Member 1, Intestinal folate carrier 1, Placental folate transporter, Reduced folate carrier protein
UniProt:	Q62866
Pathways:	Dicarboxylic Acid Transport

Application Details

Application Notes:	Western blotting: 0.5-2 µg/mL, Immunohistochemistry: 5-20 µg/mL, Immunocytochemistry: 5-20 µg/mL, Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.