

### Datasheet for ABIN7646367

# anti-SLC2A3 antibody



_					
	W	0	rv	10	W

Quantity:	100 μL	
Target:	SLC2A3	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This SLC2A3 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP)	

#### **Product Details**

Purpose:	Polyclonal Antibody to Glucose Transporter 3 (GLUT3)	
Immunogen:	RPB635Hu01Recombinant Glucose Transporter 3 (GLUT3)	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against GLUT3. It has been selected for its ability to recognize GLUT3 in immunohistochemical staining and western blotting.	
Cross-Reactivity:	Rat	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	
Target Details		
Target:	SLC2A3	

## Target Details

Alternative Name:	GLUT3 (SLC2A3 Products)	
Background:	SLC2A3, Solute Carrier Family 2(Facilitated Glucose Transporter)Member 3, Glucose transporter type 3, brain	
UniProt:	P11169	
Pathways:	Warburg Effect	

## **Application Details**

Application Notes:	Western blotting: 0.5-2 μg/mL, Immunohistochemistry: 5-20 μg/mL, Immunocytochemistry: 5-
	20 μg/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	0.3 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.