

## Datasheet for ABIN7646424

# anti-SLC44A1 antibody



#### Overview

Quantity:	100 μL
Target:	SLC44A1
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SLC44A1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

#### **Product Details**

1 Toddet Details	
Purpose:	Polyclonal Antibody to Choline Transporter Like Protein 1 (CTL1)
Immunogen:	RPB858Hu01Recombinant Choline Transporter Like Protein 1 (CTL1)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against CTL1. It has been selected for its ability to recognize CTL1 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse, Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	
Target:	SLC44A1

### **Target Details**

Alternative Name:	CTL1 (SLC44A1 Products)
Background:	CD92, CDw92, SLC44A1, CHTL1, SLC44-A1, Solute Carrier Family 44 Member 1
UniProt:	Q8WWI5

## **Application Details**

Application Notes:	Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100 Immunocytochemistry: 5-20 μg/mL,1:25-100 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	0.77 mg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.