

Datasheet for ABIN7646464
anti-SLC9A1 antibody



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	SLC9A1
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This SLC9A1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Monoclonal Antibody to Sodium/Hydrogen Exchanger 1 (NHE1)
Specificity:	The antibody is a mouse monoclonal antibody raised against NHE1. It has been selected for its ability to recognize NHE1 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	SLC9A1
Alternative Name:	NHE1 (SLC9A1 Products)
Background:	SLC9A1, APNH, Solute Carrier Family 9 Member 1, Sodium/Hydrogen Exchanger, Antiporter, Na ⁺ /H ⁺ , Amiloride Sensitive, Na(+)/H(+) antiporter, amiloride-sensitive
UniProt:	P19634

Target Details

Pathways: [Glycosaminoglycan Metabolic Process](#), [Proton Transport](#)

Application Details

Application Notes:	Western blotting: 0.2-2 µg/mL, 1:500-5000 Immunohistochemistry: 5-20 µg/mL, 1:50-200 Immunocytochemistry: 5-20 µg/mL, 1:50-200 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.