

Datasheet for ABIN7646657

anti-SDHA antibody



_					
	W	0	rv	10	W

Quantity:	100 μL	
Target:	SDHA	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This SDHA antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

Product Details

Target:

Alternative Name:

SDHA

SDHA (SDHA Products)

Purpose:	Polyclonal Antibody to Succinate Dehydrogenase Complex Subunit A (SDHA)	
Immunogen:	RPJ784Mu02Recombinant Succinate Dehydrogenase Complex Subunit A (SDHA)	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against SDHA. It has been selected for its ability to recognize SDHA in immunohistochemical staining and western blotting.	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	
Target Details		

Target Details

Background:	SDH2, SDHF, FP, Flavoprotein, Flavoprotein subunit of complex II, Succinate dehydrogenase		
	[ubiquinone] flavoprotein subunit, mitochondrial		
UniProt:	Q8K2B3		
Application Details			
Application Notes:	Western blotting: 0.5-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunocytochemistry: 5-20 μg/mL,Optimal working dilutions must be determined by end user.		
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	500 μg/mL		
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.		
Storage:	4 °C,-20 °C		
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.		