

Datasheet for ABIN7646728

anti-SOCS2 antibody



| _ | | | | | | |
|---|-------|-------------|----|----|-------------|-----|
| | V | \triangle | r۱ | /1 | \triangle | Λ/ |
| | ' V ' | | ΙV | | | v v |

Alternative Name:

| Quantity: | 100 μL | | |
|-------------------|--|--|--|
| Target: | SOCS2 | | |
| Reactivity: | Rat | | |
| Host: | Rabbit | | |
| Clonality: | Polyclonal | | |
| Conjugate: | This SOCS2 antibody is un-conjugated | | |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC) | | |
| Product Details | | | |
| Purpose: | Polyclonal Antibody to Suppressors Of Cytokine Signaling 2 (SOCS2) | | |
| Immunogen: | RPH154Ra01Recombinant Suppressors Of Cytokine Signaling 2 (SOCS2) | | |
| Isotype: | IgG | | |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against SOCS2. It has been selected for its ability to recognize SOCS2 in immunohistochemical staining and western blotting. | | |
| Cross-Reactivity: | Human, Pig | | |
| Purification: | Antigen-specific affinity chromatography followed by Protein A affinity chromatography | | |
| Target Details | | | |
| Target: | SOCS2 | | |
| | | | |

SOCS2 (SOCS2 Products)

Target Details

| Target Details | | | |
|---------------------|---|--|--|
| Background: | CIS2, Cish2, SOCS-2, SSI-2, SSI2, STATI2, Cytokine-inducible SH2 protein 2, STAT-induced STA inhibitor 2 | | |
| UniProt: | 088582 | | |
| Pathways: | JAK-STAT Signaling, Response to Growth Hormone Stimulus | | |
| Application Details | | | |
| Application Notes: | Western blotting: 0.01-2 μ g/mL,Immunohistochemistry: 5-20 μ g/mL,Optimal working dilutions must be determined by end user. | | |
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. | | |
| Restrictions: | For Research Use only | | |
| Handling | | | |
| Format: | Liquid | | |
| Concentration: | 0.5 mg/mL | | |
| Buffer: | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. | | |
| Preservative: | ProClin | | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. | | |
| Storage: | 4 °C,-20 °C | | |
| Storage Comment: | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without | | |

detectable loss of activity. Avoid repeated freeze-thaw cycles.