

Datasheet for ABIN7646850

anti-TAC1 antibody



Overview

Quantity:	100 μL
Target:	TAC1
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This TAC1 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Western Blotting (WB), Immunocytochemistry (ICC), Immunoprecipitation (IP)

Product Details

Purpose:	Monoclonal Antibody to Tachykinin, Precursor 1 (TAC1)
Specificity:	The antibody is a mouse monoclonal antibody raised against TAC1. It has been selected for its
	ability to recognize TAC1 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	TAC1
Alternative Name:	TAC1 (TAC1 Products)
Background:	NK2, NKNA, NKA, TAC2, •,PPT, Protachykinin-1, Substance P, Neurokinin A, Neuromedin L, Substance K, Neuropeptide K, Neuropeptide gamma, C-terminal-flanking peptide
UniProt:	P20366

Application Details

Application Notes:	Western blotting: $0.2-2~\mu g/m L$, $1:500-5000~lmmunohistochemistry: 5-20~\mu g/m L, 1:50-200~lmmunocytochemistry: 5-20~\mu g/m L, 1:50-200~Optimal~working~dilutions~must~be~determined~by~end~user.$
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.