

Datasheet for ABIN7646854 **anti-TAF13 antibody**



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	TAF13
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TAF13 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Western Blotting (WB), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to TATA Box Binding Protein Associated Factor 13 (TAF13)
Immunogen:	RPE563Hu01Recombinant TATA Box Binding Protein Associated Factor 13 (TAF13)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against TAF13. It has been selected for its ability to recognize TAF13 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	TAF13
Alternative Name:	TAF13 (TAF13 Products)

Target Details

Background: TAF2K, TAFII18, Transcription Initiation Factor TFIID Subunit 13, Transcription initiation factor TFIID 18 kDa subunit

UniProt: [Q15543](#)

Application Details

Application Notes: Western blotting: 0.5-2 µg/mL, Immunohistochemistry: 5-20 µg/mL, Immunocytochemistry: 5-20 µg/mL, Optimal working dilutions must be determined by end user.

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 mg/mL

Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.