

Datasheet for ABIN7646952

anti-TXNIP antibody



Overview

Quantity:	100 μL
Target:	TXNIP
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TXNIP antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Target:

Alternative Name:

Background:

TXNIP

TXNIP (TXNIP Products)

Purpose:	Polyclonal Antibody to Thioredoxin-interacting Protein (TXNIP)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against TXNIP. It has been selected for its ability to recognize TXNIP in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	

1, Upregulated By 1,25-Dihydroxyvitamin D-3

TBP2, HHCPA78, THIF, VDUP1, Thioredoxin Binding Protein 2, Vitamin D3-Upregulated Protein

Target Details

UniProt:	Q8BG60
Pathways:	Protein targeting to Nucleus, Platelet-derived growth Factor Receptor Signaling, Inflammasome
Application Details	
Application Notes:	Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100 Immunocytochemistry: 5-20 μg/mL,1:25-100 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	500 μg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.