

## Datasheet for ABIN7647324 **anti-TRRAP antibody**



[Go to Product page](#)

### Overview

Quantity:	100 µL
Target:	TRRAP
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRRAP antibody is un-conjugated
Application:	Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

### Product Details

Purpose:	Polyclonal Antibody to Transformation/Transcription Domain Associated Protein (TRRAP)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against TRRAP. It has been selected for its ability to recognize TRRAP in immunohistochemical staining and western blotting.
Cross-Reactivity:	Pig
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

### Target Details

Target:	TRRAP
Alternative Name:	TRRAP ( <a href="#">TRRAP Products</a> )

## Target Details

Background: STAF40, TR-AP, Tra1, 350/400 kDa PCAF-associated factor

UniProt: [Q9Y4A5](#)

## Application Details

Application Notes: Western blotting: 0.2-2 µg/mL, 1:250-2500 Immunohistochemistry: 5-20 µg/mL, 1:25-100  
Immunocytochemistry: 5-20 µg/mL, 1:25-100 Optimal working dilutions must be determined by end user.

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: 500 µg/mL

Buffer: PBS, pH 7.4, containing 0.01 % SKL, 1 mM DTT, 5 % Trehalose and Proclin300.

Preservative: Dithiothreitol (DTT), ProClin

Precaution of Use: This product contains ProClin and Dithiothreitol (DTT): POISONOUS AND HAZARDOUS SUBSTANCES which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.