

## Datasheet for ABIN7647345 **anti-TGFB1 antibody**



[Go to Product page](#)

### Overview

|              |                                                                                                           |
|--------------|-----------------------------------------------------------------------------------------------------------|
| Quantity:    | 100 µL                                                                                                    |
| Target:      | TGFB1                                                                                                     |
| Reactivity:  | Zebrafish (Danio rerio)                                                                                   |
| Host:        | Rabbit                                                                                                    |
| Clonality:   | Polyclonal                                                                                                |
| Conjugate:   | This TGFB1 antibody is un-conjugated                                                                      |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP),<br>Immunocytochemistry (ICC) |

### Product Details

|               |                                                                                                                                                                                  |
|---------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Purpose:      | Polyclonal Antibody to Transforming Growth Factor Beta 1 (TGFb1)                                                                                                                 |
| Immunogen:    | RPA124Fi01Recombinant Transforming Growth Factor Beta 1 (TGFb1)                                                                                                                  |
| Isotype:      | IgG                                                                                                                                                                              |
| Specificity:  | The antibody is a rabbit polyclonal antibody raised against TGFb1. It has been selected for its ability to recognize TGFb1 in immunohistochemical staining and western blotting. |
| Purification: | Antigen-specific affinity chromatography followed by Protein A affinity chromatography                                                                                           |

### Target Details

|                   |                                          |
|-------------------|------------------------------------------|
| Target:           | TGFB1                                    |
| Alternative Name: | TGFb1 ( <a href="#">TGFB1 Products</a> ) |

## Target Details

|             |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
|-------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Background: | TGF-B1, CED, DPD1, LAP, Camurati-Engelmann Disease, Latency-associated peptide                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| UniProt:    | <a href="#">Q7ZZU7</a>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Pathways:   | <a href="#">EGFR Signaling Pathway</a> , <a href="#">Dopaminergic Neurogenesis</a> , <a href="#">Cellular Response to Molecule of Bacterial Origin</a> , <a href="#">Glycosaminoglycan Metabolic Process</a> , <a href="#">Regulation of Leukocyte Mediated Immunity</a> , <a href="#">Regulation of Muscle Cell Differentiation</a> , <a href="#">Positive Regulation of Immune Effector Process</a> , <a href="#">Cell-Cell Junction Organization</a> , <a href="#">Production of Molecular Mediator of Immune Response</a> , <a href="#">Ribonucleoside Biosynthetic Process</a> , <a href="#">Skeletal Muscle Fiber Development</a> , <a href="#">Regulation of Carbohydrate Metabolic Process</a> , <a href="#">Protein targeting to Nucleus</a> , <a href="#">Autophagy</a> , <a href="#">Cancer Immune Checkpoints</a> |

## Application Details

|                    |                                                                                                                                                                                                                                                                                                                                     |
|--------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Application Notes: | Western blotting: 0.5-2 µg/mL, Immunohistochemistry: 5-20 µg/mL, Immunocytochemistry: 5-20 µg/mL, Optimal working dilutions must be determined by end user.                                                                                                                                                                         |
| Comment:           | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. |
| Restrictions:      | For Research Use only                                                                                                                                                                                                                                                                                                               |

## Handling

|                    |                                                                                                                                                                 |
|--------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Format:            | Liquid                                                                                                                                                          |
| Concentration:     | 0.5 mg/mL                                                                                                                                                       |
| Buffer:            | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.                                                                                                |
| Preservative:      | ProClin                                                                                                                                                         |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.                                               |
| Storage:           | 4 °C, -20 °C                                                                                                                                                    |
| Storage Comment:   | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles. |