

Datasheet for ABIN7647425
anti-Transglutaminase 2 antibody



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	Transglutaminase 2 (TGM2)
Reactivity:	Mouse
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Transglutaminase 2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Monoclonal Antibody to Transglutaminase 2 (TGM2)
Immunogen:	RPB830Mu01Recombinant Transglutaminase 2 (TGM2)
Clone:	C2
Specificity:	The antibody is a mouse monoclonal antibody raised against TGM2. It has been selected for its ability to recognize TGM2 in immunohistochemical staining and western blotting.
Purification:	Protein A + Protein G affinity chromatography

Target Details

Target:	Transglutaminase 2 (TGM2)
Alternative Name:	TGM2 (TGM2 Products)

Target Details

Background:	TTG, TG2, C Polypeptide,Protein-Glutamine-Gamma-Glutamyltransferase, Transglutaminase 2, Tissue, Transglutaminase C, Transglutaminase H
UniProt:	P21981
Pathways:	Tube Formation , Thromboxane A2 Receptor Signaling

Application Details

Application Notes:	Western blotting: 0.5-2 µg/mL,Immunohistochemistry: 5-20 µg/mL,Immunocytochemistry: 5-20 µg/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.