

## Datasheet for ABIN7647482 **anti-TMEFF1 antibody**



[Go to Product page](#)

### Overview

Quantity:	100 µL
Target:	TMEFF1
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TMEFF1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

### Product Details

Purpose:	Polyclonal Antibody to Tomoregulin 1 (TR1)
Immunogen:	RPM377Mu01Recombinant Tomoregulin 1 (TR1)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against TR1. It has been selected for its ability to recognize TR1 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

### Target Details

Target:	TMEFF1
Alternative Name:	Tomoregulin 1 ( <a href="#">TMEFF1 Products</a> )

## Target Details

Background: TMEFF1, CT120.1, C9orf2, Transmembrane Protein With EGF-Like And Two Follistatin-Like Domains 1, Cancer/Testis Antigen Family 120,Member 1

UniProt: [Q6PFE7](#)

## Application Details

Application Notes: Western blotting: 0.01-2 µg/mL,Immunohistochemistry: 5-20 µg/mL,Immunocytochemistry: 5-20 µg/mL,Optimal working dilutions must be determined by end user.

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: 0.5 mg/mL

Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-20 °C

Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.