

### Datasheet for ABIN7647863

# anti-UGT8 antibody



#### Overview

Quantity:	100 μL
Target:	UGT8
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This UGT8 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

#### **Product Details**

Floduct Details	
Purpose:	Polyclonal Antibody to UDP Glycosyltransferase 8 (UGT8)
Immunogen:	RPG919Hu01Recombinant UDP Glycosyltransferase 8 (UGT8)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against UGT8. It has been selected for its ability to recognize UGT8 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse, Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	
Target:	UGT8

## **Target Details**

Alternative Name:	UGT8 (UGT8 Products)
Background:	CGT, Cerebroside synthase, UDP-Galactose Ceramide Galactosyltransferase, 2- Hydroxyacylsphingosine 1-Beta-Galactosyltransferase
UniProt:	Q16880
Pathways:	Cell-Cell Junction Organization
Application Details	

Restrictions:	For Research Use only
	date under appropriate storage condition.
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	20 μg/mL,Optimal working dilutions must be determined by end user.
Application Notes:	Western blotting: 0.5-2 $\mu$ g/mL,Immunohistochemistry: 5-20 $\mu$ g/mL,Immunocytochemistry: 5-

## Handling

Format:	Liquid
Concentration:	0.23 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.