

Datasheet for ABIN7647964 **anti-VTCN1 antibody**



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	VTCN1
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This VTCN1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to V-Set Domain Containing T-Cell Activation Inhibitor 1 (VTCN1)
Immunogen:	RPK032Hu03Recombinant VSet Domain Containing TCell Activation Inhibitor 1 (VTCN1)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against VTCN1. It has been selected for its ability to recognize VTCN1 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	VTCN1
Alternative Name:	VTCN1 (VTCN1 Products)

Target Details

Background: B7-H4, B7H4, B7S1, B7X, B7h.5, VCTN1, T-cell costimulatory molecule B7x, Immune costimulatory protein B7-H4

UniProt: [Q7Z7D3](#)

Application Details

Application Notes: Western blotting: 0.01-2 µg/mL, Immunohistochemistry: 5-20 µg/mL, Immunocytochemistry: 5-20 µg/mL, Optimal working dilutions must be determined by end user.

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 0.5 mg/mL

Buffer: 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.