

Datasheet for ABIN7648050 anti-Vip antibody



| Overview | |
|-------------------|--|
| Quantity: | 100 μL |
| Target: | Vip |
| Reactivity: | Cow |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This Vip antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |
| Product Details | |
| Purpose: | Polyclonal Antibody to Vasoactive Intestinal Peptide (VIP) |
| Immunogen: | RPA380Bo01Recombinant Vasoactive Intestinal Peptide (VIP) |
| Isotype: | IgG |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against VIP. It has been selected for its ability to recognize VIP in immunohistochemical staining and western blotting. |
| Purification: | Antigen-specific affinity chromatography followed by Protein A affinity chromatography |
| Target Details | |
| Target: | Vip |
| Alternative Name: | Vasoactive Intestinal Peptide (Vip Products) |

Target Details

Storage Comment:

| Target Details | |
|---------------------|---|
| Background: | PHM27, Intestinal peptide PHV-42, Peptide histidine valine 42, Peptide histidine methioninamide 27 |
| UniProt: | P81401 |
| Pathways: | Hormone Activity, cAMP Metabolic Process |
| Application Details | |
| Application Notes: | Western blotting: 0.01-2 μ g/mL,Immunohistochemistry: 5-20 μ g/mL,Immunocytochemistry: 5-20 μ g/mL,Optimal working dilutions must be determined by end user. |
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 0.44 mg/mL |
| Buffer: | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| | |

detectable loss of activity. Avoid repeated freeze-thaw cycles.

Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without