

Datasheet for ABIN7648228

anti-WISP1 antibody



_					
	W	0	rv	10	W

Quantity:	100 μL	
Target:	WISP1	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This WISP1 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

Product Details

Target:

Alternative Name:

WISP1

WISP1 (WISP1 Products)

Purpose:	Polyclonal Antibody to WNT1 Inducible Signaling Pathway Protein 1 (WISP1)
Immunogen:	RPG895Hu01Recombinant WNT1 Inducible Signaling Pathway Protein 1 (WISP1)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against WISP1. It has been selected for its ability to recognize WISP1 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	

Target Details

Background:	CCN4, WISP1c, WISP1i, WISP1tc, CCN family member 4, Wnt-1-induced secreted protein	
UniProt:	095388	
Pathways:	WNT Signaling, Growth Factor Binding	

Application Details

Application Notes:	Western blotting: 0.01-2 μ g/mL,Immunohistochemistry: 5-20 μ g/mL,Immunocytochemistry: 5-	
	20 μg/mL,Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
	date under appropriate storage condition.	
Restrictions:	For Research Use only	

Handling

Format:	Liquid	
Concentration:	0.5 mg/mL	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.	