

### Datasheet for ABIN7648273

# anti-XPA antibody



#### Overview

Quantity:	100 μL
Target:	XPA
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This XPA antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

### **Product Details**

Alternative Name:

Product Details	
Purpose:	Polyclonal Antibody to Xeroderma Pigmentosum, Complementation Group A (XPA)
Immunogen:	RPE744Hu01Recombinant Xeroderma Pigmentosum, Complementation Group A (XPA)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against XPA. It has been selected for its ability to recognize XPA in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	
Target:	XPA

Xeroderma Pigmentosum, Complementation Group A (XPA Products)

## **Target Details**

Background:	XP-A, XPAC, XP1, DNA repair protein complementing XP-A cells
UniProt:	P23025
Pathways:	DNA Damage Repair

## **Application Details**

Application Notes:	Western blotting: 0.01-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunocytochemistry: 5-
	20 μg/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only

# Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.