

Datasheet for ABIN7648278

anti-FXYD2 antibody



Overview

Quantity:	100 μL
Target:	FXYD2
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This FXYD2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP)

Product Details

Target:

- Troduct Details	
Purpose:	Monoclonal Antibody to Hypomagnesemia 2, Renal (HOMG2)
Immunogen:	CPL527Hu11BSA Conjugated Hypomagnesemia 2, Renal (HOMG2)
Clone:	C1
Specificity:	The antibody is a mouse monoclonal antibody raised against HOMG2. It has been selected for its ability to recognize HOMG2 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Pig
Purification:	Protein A + Protein G affinity chromatography
Target Details	

FXYD2

Target Details

rarget Details	
Alternative Name:	HOMG2 (FXYD2 Products)
Background:	ATP1G1, FXYD2, ATP1C, Sodium pump gamma chain, FXYD Domain Containing Ion Transport Regulator 2, Sodium/Potassium-Transporting ATPase Gamma Chain
UniProt:	P54710
Pathways:	Thyroid Hormone Synthesis
Application Details	
Application Notes:	Western blotting: 0.2-2 μ g/mL,1:500-5000 Immunohistochemistry: 5-20 μ g/mL,1:50-200 Immunocytochemistry: 5-20 μ g/mL,1:50-200 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin, Sodium azide
Precaution of Use:	This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANCES which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.