

Datasheet for ABIN7648602 **anti-FCAR antibody**



[Go to Product page](#)

Overview

| | |
|--------------|-------------------------------------|
| Quantity: | 100 tests |
| Target: | FCAR |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This FCAR antibody is un-conjugated |
| Application: | Flow Cytometry (FACS) |

Product Details

| | |
|---------------|--|
| Purpose: | Anti-Fc Fragment Of IgA Receptor (FcaR) Polyclonal Antibody |
| Isotype: | IgG |
| Purification: | Antigen-specific affinity chromatography followed by Protein A affinity chromatography |

Target Details

| | |
|-------------------|--|
| Target: | FCAR |
| Alternative Name: | Fc Fragment Of IgA Receptor (FCAR Products) |
| Background: | CD89, FcaRI, Receptor I For The Fc Region Of Immunoglobulin A, IgA Fc receptor, Immunoglobulin alpha Fc receptor |
| UniProt: | P24071 |

Application Details

Application Notes: For flow cytometry, the suggested use of this reagent is 1-5 μL per 10^6 cells in 100 μL volume. Optimal working dilutions could be determined by end user.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.