

## Datasheet for ABIN7648672 **anti-ICOS antibody (FITC)**

[Go to Product page](#)

### Overview

|              |                                          |
|--------------|------------------------------------------|
| Quantity:    | 100 tests                                |
| Target:      | ICOS                                     |
| Reactivity:  | Human                                    |
| Host:        | HEK-293F Cells                           |
| Clonality:   | Polyclonal                               |
| Conjugate:   | This ICOS antibody is conjugated to FITC |
| Application: | Flow Cytometry (FACS)                    |

### Product Details

|               |                                                                                        |
|---------------|----------------------------------------------------------------------------------------|
| Purpose:      | FITC-Linked Anti-Inducible T-Cell Co Stimulator (ICOS) Polyclonal Antibody             |
| Isotype:      | IgG                                                                                    |
| Purification: | Antigen-specific affinity chromatography followed by Protein A affinity chromatography |

### Target Details

|                   |                                                                                                       |
|-------------------|-------------------------------------------------------------------------------------------------------|
| Target:           | ICOS                                                                                                  |
| Alternative Name: | Inducible T-Cell Co Stimulator ( <a href="#">ICOS Products</a> )                                      |
| Background:       | CD278, AILIM, Inducible T-Cell Costimulator, Activation-Inducible Lymphocyte Immunomediatory Molecule |
| UniProt:          | <a href="#">Q9Y6W8</a>                                                                                |
| Pathways:         | <a href="#">Cancer Immune Checkpoints</a>                                                             |

Application Details

Application Notes: For flow cytometry, the suggested use of this reagent is 1-5 µL per 10<sup>6</sup> cells in 100 µL volume.  
Optimal working dilutions could be determined by end user.

Restrictions: For Research Use only

Handling

|                    |                                                                                                                   |
|--------------------|-------------------------------------------------------------------------------------------------------------------|
| Format:            | Liquid                                                                                                            |
| Buffer:            | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.                                                  |
| Preservative:      | ProClin                                                                                                           |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage:           | 4 °C                                                                                                              |
| Storage Comment:   | Store at 4°C, protected from prolonged exposure to light.                                                         |