

Datasheet for ABIN7651616

Recombinant anti-CD45RB antibody



[Go to Product page](#)

Overview

Quantity:	50 µL
Target:	CD45RB
Reactivity:	Human
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This CD45RB antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunofluorescence (IF), Immunohistochemistry (IHC), Western Blotting (WB), Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections) (IHC (fp))

Product Details

Purpose:	CD45RB (B-Cell Marker) (PTPRC/2877R)
Immunogen:	Recombinant full-length human PTPRC protein
Clone:	PTPRC-2877R
Isotype:	IgG
Characteristics:	CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these

Product Details

isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45RB is expressed on mature B-lymphocytes and the majority of lymphomas and leukemias of B-cell origin. Primary antibodies are available purified, or with a selection of fluorescent CF® Dyes and other labels. CF® Dyes offer exceptional brightness and photostability. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Target Details

Target:	CD45RB
Alternative Name:	CD45RB
Background:	Synonyms: B220, CD45R, GP180, Leukocyte common antigen (LCA), Loc, Ly-5, Lyt-4, Protein tyrosine phosphatase receptor type C (PTPRC), Receptor-type tyrosine-protein phosphatase C, T200 glycoprotein Gene Symbol: PTPRC Tissue Expression: B-cells
Molecular Weight:	180-220 kDa
Gene ID:	5788
UniProt:	P08575

Application Details

Application Notes:	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Does not react with rat or rhesus monkey. Immunohistology (formalin): 0.5-1.0 µg/mL for 30 minutes at RT. Western Blot 0.5-2 µg/mL. Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined by user
Comment:	Positive Control: Raji, Ramos, U-698, or GA-10 cells. Tonsil
Restrictions:	For Research Use only

Handling

Format:	Liquid
---------	--------

Handling

Concentration:	1 mg/mL
Buffer:	PBS, no BSA, no azide
Preservative:	Without preservative
Storage:	-20 °C
Storage Comment:	Stable at room temperature or 37°C for 7 days.
	Store at -20 °C. Protect fluorescent conjugates from light
Expiry Date:	24 months