

Datasheet for ABIN7659758

anti-L-Selectin antibody (Biotin)



Overview

Quantity:	100 μL
Target:	L-Selectin (SELL)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This L-Selectin antibody is conjugated to Biotin
Application:	Please inquire

Product Details

Purpose:	CD62L (L-Selectin)(CD62L/1588), Biotin conjugate
Immunogen:	MAb raised against supernatant from phorbol myristic acid activated human peripheral blood leukocytes.
Clone:	CD62L-1588
Isotype:	IgG1, kappa
Characteristics:	Salactine also decignated CD62 antigons comprise a family of earbohydrate-hinding proteins

Characteristics:

Selectins, also designated CD62 antigens, comprise a family of carbohydrate-binding proteins involved in mediating cellular interactions with leukocytes. L-Selectin (also designated LECAM-1 or CD62L) is expressed on the majority of B and naive T cells and on most monocytes, neutrophils and eosinophils. L-Selectin interacts with specific carbohydrates expressed by activated endothelial cells. P-Selectin (also designated GMP-140 or CD62P), expressed on activated platelets and endothelial cells, and E-Selectin (also designated ELMA-1 or CD62E), expressed on endothelial cells, exhibit overlapping ligand specificities. Both recognize sialyl-Le

(x) as a ligand and bind to specific carbohydrates on neutrophils and monocytes. Primary antibodies are available purified, or with a selection of fluorescent CF® Dyes and other labels. CF® Dyes offer exceptional brightness and photostability. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Target Details

Handling

Format:

rarget Details	
Target:	L-Selectin (SELL)
Alternative Name:	CD62L
Background:	Synonyms: A.11, AI528707, CD62 antigen ligand, CD62L, gp90-MEL, IgA nephropathy,
	susceptibility to, included, L Selectin, L-selectin, LAM1, LECAM1, LEU8, Leukocyte surface
	antigen Leu-8, Leukocyte-endothelial cell adhesion molecule 1, Lnhr, LSEL, Ly-22, LYAM1,
	Lymph node homing receptor, Lymphocyte antigen 22, Lymphocyte surface MEL-14 antigen,
	Pln homing receptor, PLNHR, SELL, TQ1
	Gene Symbol: SELL
	Tissue Expression: Eosinophils Lymphocytes Monocytes/macrophages Neutrophils
Molecular Weight:	65-75 kDa
Gene ID:	6402
UniProt:	P14151
Application Details	
Application Notes:	For coating for ELISA, order Ab without BSA. Higher concentration may be required for direct
	detection using primary antibody conjugates than for indirect detection with secondary
	antibody. Optimal dilution and staining procedure for a specific application should be
	determined by user. Recommended starting concentrations for titration are 1-2 µg/mL for mos
	applications, or 1 μg/million cells/100 μLfor flow cytometry
Comment:	Positive Control: HPBL or Jurkat whole cell lysate.
Restrictions:	For Research Use only

Liquid

Handling

Concentration:	0.1 mg/mL
Buffer:	PBS, 0.1 % BSA, 0.05 % azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Stable at room temperature or 37°C for 7 days. Store at 2 to 8°C. Protect fluorescent conjugates from light
Expiry Date:	24 months