



[Go to Product page](#)

Datasheet for ABIN775705

anti-CD28 antibody

4 Images

5 Publications

Overview

Quantity:	0.1 mg
Target:	CD28
Reactivity:	Human, Non-Human Primate
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD28 antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunoprecipitation (IP), Western Blotting (WB), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	DC28.1.3.3 murine T cell hybridoma transfected with human CD28 cDNA
Clone:	CD28-2
Isotype:	IgG1 kappa
Specificity:	The mouse monoclonal antibody CD28.2 recognizes an extracellular epitope of CD28, a disulfide-linked homodimeric type I glycoprotein (monomer of Mw 44 kDa) which is a critical costimulatory receptor of T cells.
Cross-Reactivity (Details):	Human, Non-Human Primates
Purification:	Purified by protein-A affinity chromatography.
Purity:	> 95 % (by SDS-PAGE)

Target Details

Target:	CD28
Alternative Name:	CD28 (CD28 Products)
Background:	CD28 Molecule,CD28 is the critical T cell costimulatory receptor which provides to the cell the important second activation signal by binding CD80 and CD86 that are expressed by antigen presenting cells. Besides its costimulation role CD28 functions in preventing T cells from anergic hyporesponsive state or from undergoing premature apoptotic cell death. CD28 is also expressed on human fetal NK cells and some NK cell lines, whereas on murine NK cells the CD28 expression is much broader.,TP44
Gene ID:	940
UniProt:	P10747
Pathways:	TCR Signaling , Fc-epsilon Receptor Signaling Pathway , EGFR Signaling Pathway , Regulation of Leukocyte Mediated Immunity , Positive Regulation of Immune Effector Process , Production of Molecular Mediator of Immune Response

Application Details

Application Notes:	Flow cytometry: Recommended dilution: 1-12 µg/mL
Restrictions:	For Research Use only

Handling

Concentration:	1 mg/mL
Buffer:	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Do not freeze.
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Do not freeze.

Publications

Product cited in:	Jeong, Qiao, Nascimbeni, Hu, Rehermann, Murthy, Liang: "Immunization with hepatitis C virus-
-------------------	--

like particles induces humoral and cellular immune responses in nonhuman primates." in:

Journal of virology, Vol. 78, Issue 13, pp. 6995-7003, (2004) ([PubMed](#)).


Scharschmidt, Wegener, Heissmeyer, Rao, Krappmann: "Degradation of Bcl10 induced by T-cell activation negatively regulates NF-kappa B signaling." in: **Molecular and cellular biology**, Vol. 24, Issue 9, pp. 3860-73, (2004) ([PubMed](#)).

Marti, Krause, Post, Lyddane, Dupont, Sadelain, King: "Negative-feedback regulation of CD28 costimulation by a novel mitogen-activated protein kinase phosphatase, MKP6." in: **Journal of immunology (Baltimore, Md. : 1950)**, Vol. 166, Issue 1, pp. 197-206, (2001) ([PubMed](#)).

Galea-Lauri, Darling, Gan, Krivochtchapov, Kuiper, Gäken, Souberbielle, Farzaneh: "Expression of a variant of CD28 on a subpopulation of human NK cells: implications for B7-mediated stimulation of NK cells." in: **Journal of immunology (Baltimore, Md. : 1950)**, Vol. 163, Issue 1, pp. 62-70, (1999) ([PubMed](#)).


Tazi, Moreau, Bergeron, Dominique, Hance, Soler: "Evidence that Langerhans cells in adult pulmonary Langerhans cell histiocytosis are mature dendritic cells: importance of the cytokine microenvironment." in: **Journal of immunology (Baltimore, Md. : 1950)**, Vol. 163, Issue 6, pp. 3511-5, (1999) ([PubMed](#)).

Images




Western Blotting

Image 1. Western blotting analysis of human CD28 using mouse monoclonal antibody CD28.2 on lysates of Jurkat cells and THP1 cells (negative control) under reducing and non-reducing conditions. Nitrocellulose membrane was probed with 2 µg/mL of mouse monoclonal antibody followed by IRDye800-conjugated anti-mouse secondary antibody. CD28 was detected around 35-40 kDa.



Flow Cytometry

Image 2. Surface staining of human peripheral blood leukocytes with anti-human CD28 (CD28.2) purified.



Flow Cytometry

Image 3. Separation of human CD28 positive lymphocytes (red-filled) from neutrophil granulocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD28 (CD28.2) purified antibody (concentration in sample 1,6 μ g/mL) GAM APC.

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN775705.