



[Go to Product page](#)

Datasheet for ABIN775708  
**anti-Lambda-IgLC antibody**

[2 Images](#)

[3 Publications](#)

### Overview

Quantity:	0.1 mg
Target:	Lambda-IgLC
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Lambda-IgLC antibody is un-conjugated
Application:	Flow Cytometry (FACS)

### Product Details

Immunogen:	Human IgA1 lambda myeloma protein
Clone:	1-155-2
Isotype:	IgG1 kappa
Specificity:	The mouse monoclonal antibody 1-155-2 recognizes lambda light chains (22.5 kDa) of human immunoglobulin.
Cross-Reactivity (Details):	Human
Purification:	Purified by protein-A affinity chromatography.
Purity:	> 95 % (by SDS-PAGE)

### Target Details

Target:	Lambda-IgLC
---------	-------------

## Target Details

---

Alternative Name:	Lambda light chains ( <a href="#">Lambda-IgLC Products</a> )
Background:	Immunoglobulin classes share the same basic four polypeptide chain structure of two heavy chains (five heavy chains types) and two light chains (kappa, lambda, both having a molecular weight of 22.5 kDa). Kappa and lambda consist of a variable region and a constant region and can easily be differentiated by the antigenic properties of the constant region. The ratio of kappa to lambda is 70:30.,Immunoglobulin lambda, Igl
Molecular Weight:	22.5 kDa

## Application Details

---

Application Notes:	Flow cytometry: Recommended dilution: 1-4 µg/mL. Extracellular and intracellular staining.
Restrictions:	For Research Use only

## Handling

---

Concentration:	1 mg/mL
Buffer:	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	<b>Do not freeze.</b>
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Do not freeze.

## Publications


---

Product cited in: Tierens, Holte, Warsame, Ikonomou, Wang, Chan, Delabie: "Low levels of monoclonal small B cells in the bone marrow of patients with diffuse large B-cell lymphoma of activated B-cell type but not of germinal center B-cell type." in: **Haematologica**, Vol. 95, Issue 8, pp. 1334-41, (2010) ([PubMed](#)).

Bahler, Pindzola, Swerdlow: "Splenic marginal zone lymphomas appear to originate from different B cell types." in: **The American journal of pathology**, Vol. 161, Issue 1, pp. 81-8, (2002) ([PubMed](#)).


Kubagawa, Gathings, Levitt, Kearney, Cooper: "Immunoglobulin isotype expression of normal pre-B cells as determined by immunofluorescence." in: **Journal of clinical immunology**, Vol. 2, Issue 4, pp. 264-9, (1983) ([PubMed](#)).

Images



Flow Cytometry

**Image 1.** Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human Ig Lambda Light Chain (1-155-2) purified antibody (concentration in sample 4 µg/mL, GAM APC).



Flow Cytometry

**Image 2.** Separation of human Ig Lambda light chain positive lymphocytes (red-filled) from Ig Lambda light chain negative lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human Ig Lambda Light Chain (1-155-2) purified antibody (concentration in sample 4 µg/mL, GAM APC).