

Datasheet for ABIN790403
anti-KRT20 antibody (FITC)



[Go to Product page](#)

2 Images

Overview

Quantity:	50 µL
Target:	KRT20
Reactivity:	Human, Rat, Pig
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This KRT20 antibody is conjugated to FITC
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Flow Cytometry (FACS)

Product Details

Brand:	IHC-plus™
Immunogen:	Electrophoretically purified keratin K20 from human intestinal mucosa
Clone:	IT-Ks 20-10
Isotype:	IgG1
Specificity:	Human, pig, and rat. Polypeptide reacting: Protein IT Reactivities on cultured Cell lines that have been tested so far are HT-29, LoVo, DLD-1, SW 1116 CaCo-2, RT-4.
Purification:	Affinity purified

Target Details

Target:	KRT20
---------	-------

Target Details


Alternative Name:	KRT20 / CK20 / Cytokeratin 20 (KRT20 Products)
Background:	Name/Gene ID: KRT20 Family: Keratin Synonyms: KRT20, Cytokeratin-20, Cytokeratin 20, CK20, K20, Keratin 20, Keratin-20, KRT21, CK-20, Protein IT
Gene ID:	54474
UniProt:	P35900

Application Details

Application Notes:	Approved: Flo, IHC, IHC-P (1:50) Usage: The applications listed have been tested for the unconjugated form of this product. Other forms have not been tested.
Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	0.05 % sodium azide, 0.5 % BSA
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C,-20 °C
Storage Comment:	Short term: 4°C. Long term: Store at -20°C. Avoid freeze-thaw cycles.



Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections)

Image 1. Human Small Intestine: Formalin-Fixed, Paraffin-Embedded (FFPE)



Immunohistochemistry

Image 2. Anti-KRT20 / Cytokeratin 20 antibody IHC of human small intestine. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody dilution 1:50. This image was taken for the unconjugated form of this p ...