antibodies .- online.com





anti-FXYD1 antibody (pSer83) (Biotin)



Overview	
Quantity:	100 μL
Target:	FXYD1
Binding Specificity:	pSer83
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FXYD1 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	
Immunogen:	KLH conjugated synthetic phosphopeptide derived from human FXYD1 around the phosphorylation site of Ser83
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig
Purification:	Purified by Protein A.
Target Details	
Target:	FXYD1
Alternative Name:	FXYD1 (FXYD1 Products)

Target Details

Background:	Synonyms: FXYD1 phospho S83, FXYD1 phospho Ser83, p-FXYD1 phospho S83, FXYD domain		
	containing ion transport regulator 1, Phospholemman, PLM, fxyd1, PLM_HUMAN, FXYD		
	domain-containing ion transport regulator 1.		
	Background: PLM (FXYD1)is a member of a family of small membrane proteins that share a 35-		
	amino acid signature sequence domain, beginning with the sequence PFXYD and containing 7		
	invariant and 6 highly conserved amino acids. FXYD2, also known as the gamma subunit of the		
	Na,K-ATPase, regulates the properties of that enzyme. FXYD1 (phospholemman), FXYD2 (gamma), FXYD3 (MAT-8), FXYD4 (CHIF), and FXYD5 (RIC) have been shown to induce channel activity in experimental expression systems. PLM may be phosphorylated by several kinases, including protein kinase A, protein kinase C, NIMA kinase, and myotonic dystrophy kinase. It is		
		thought to form an ion channel or regulate ion channel activity.	
		Gene ID:	5348
		Application Details	
	Application Notes:	WB 1:300-5000	
IHC-P 1:200-400			
IHC-F 1:100-500			
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C for 12 months.		
Expiry Date:	12 months		