## antibodies -online.com





## anti-FXYD1 antibody (pSer88) (Biotin)



Overview	
Quantity:	100 μL
Target:	FXYD1
Binding Specificity:	pSer88
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FXYD1 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)),
	Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	
Immunogen:	KLH conjugated synthetic phosphopeptide derived from human FXYD1 around the
	phosphorylation site of Ser88
Isotype:	IgG
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig
Purification:	Purified by Protein A.
Target Details	
Target:	FXYD1
Alternative Name:	FXYD1 (FXYD1 Products)

## **Target Details**

•		
Background:	Synonyms: FXYD1 phospho S88, FXYD1 phospho Ser88, p-FXYD1 phospho S88, FXYD domain	
	containing ion transport regulator 1, Phospholemman, PLM, fxyd1, PLM_HUMAN, FXYD	
	domain-containing ion transport regulator 1.	
	Background: PLM (FXYD1)is a member of a family of small membrane proteins that share a 35-	
	amino acid signature sequence domain, beginning with the sequence PFXYD and containing 7	
	invariant and 6 highly conserved amino acids. FXYD2, also known as the gamma subunit of the	
	Na,K-ATPase, regulates the properties of that enzyme. FXYD1 (phospholemman), FXYD2 (gamma), FXYD3 (MAT-8), FXYD4 (CHIF), and FXYD5 (RIC) have been shown to induce channel activity in experimental expression systems. PLM may be phosphorylated by several kinases,	
		including protein kinase A, protein kinase C, NIMA kinase, and myotonic dystrophy kinase. It is
		thought to form an ion channel or regulate ion channel activity.
	Gene ID:	5348
	Application Details	
Application Notes:	WB 1:300-5000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C for 12 months.	
Expiry Date:	12 months	