

Datasheet for ABIN801417

anti-IKBKG antibody (pSer43) (HRP)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	IKBKG
Binding Specificity:	pSer43
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IKBKG antibody is conjugated to HRP
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human IKBKG around the phosphorylation site of Ser43
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse,Dog,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	IKBKG
---------	-------

Target Details

Alternative Name:	IKK gamma (IKBKG Products)
Background:	Synonyms: IκB kinase associated protein 1, IκB kinase subunit gamma, Inhibitor of nuclear factor kappa B kinase subunit gamma, AMCBX1, FIP 3, FIP3, Fip3p, I kappa B kinase gamma, IκB kinase associated protein 1, IκB kinase gamma subunit, IκB kinase subunit gamma, IKBKG, IKKAP 1, IKKAP1, IKKG, IKKgamma,IKK gamma. Background: Pro inflammatory cytokines activate the transcription factor NF kappa B by stimulating the activity of a protein kinase that phosphorylates Iκappa B, an inhibitor of NF kappa B, at sites that trigger its ubiquitination and degradation. A large, cytokine responsive Iκappa B kinase (IKK) complex has been purified and the genes encoding 2 of its subunits have been cloned. These subunits, IKK alpha and Iκk beta, are protein kinases whose function is needed for NF kappa B activation by pro inflammatory stimuli. IKK is composed of similar amounts of IKK alpha, Iκk beta, which are differentially processed forms of a third subunit, IKK gamma. IKK gamma interacts preferentially with IKK beta and is required for the activation of the IKK complex.
Gene ID:	8517
Pathways:	NF-kappaB Signaling , RTK Signaling , TCR Signaling , TLR Signaling , Fc-epsilon Receptor Signaling Pathway , Activation of Innate immune Response , M Phase , Production of Molecular Mediator of Immune Response , Hepatitis C , Protein targeting to Nucleus , Toll-Like Receptors Cascades , BCR Signaling , Ubiquitin Proteasome Pathway , S100 Proteins

Application Details

Application Notes:	IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

	handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months