

Datasheet for ABIN801942

anti-MAP2K5 antibody (pSer142) (HRP)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	MAP2K5
Binding Specificity:	pSer142
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MAP2K5 antibody is conjugated to HRP
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human MEK5 around the phosphorylation site of Ser142
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	MAP2K5
Alternative Name:	MEK5 (MAP2K5 Products)

Target Details

Background:	Synonyms: MAP2K5 phospho S142, Dual specificity mitogen activated protein kinase kinase 5, MAP kinase kinase 5, MAP kinase kinase MEK5b, MAP2K5, MAPK/ERK kinase 5, MAPKK5, PRKMK5, EC 2.7.12.2, MP2K5_HUMAN, Dual specificity mitogen-activated protein kinase kinase 5, MAPKK 5, MAPK/ERK kinase 5, MEK 5. Background: MEK5 is a dual specificity protein kinase belonging to the Ser/Thr protein kinase family, (MAP kinase kinase family). It is activated by phosphorylation on Ser/Thr by MAP kinase kinases and interacts specifically with ERK5, and not with another MAP kinase like P38. This kinase specifically interacts with and activates MAPK7/ERK5. This kinase itself can be phosphorylated and activated by MAP3K3/MEKK3, as well as by atypical protein kinase C isoforms (aPKCs). It is not phosphorylated by RAFA, RAFB or RAFC and it may interact with GTPases such as CDC42. The signal cascade mediated by this kinase is involved in growth factor stimulated cell proliferation and muscle cell differentiation. MEK5 is expressed in many adult tissues and is most abundant in heart and skeletal muscle.
Gene ID:	5607
Pathways:	MAPK Signaling , Neurotrophin Signaling Pathway

Application Details

Application Notes:	IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.
Storage:	-20 °C

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months