



[Go to Product page](#)

Datasheet for ABIN863105  
**anti-Piccolo antibody**

2 Images

Overview

Quantity:	100 µg
Target:	Piccolo (PIC)
Reactivity:	Mouse
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Piccolo antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Immunofluorescence (IF)

Product Details

Immunogen:	Full length recombinant protein (GenBank ID: BC001304.1)
Clone:	6H9-B6
Isotype:	IgG2b kappa
Specificity:	Detects ~550 kDa. No cross-reactivity to Bassoon.
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Protein G Purified

Target Details

Target:	Piccolo (PIC)
Alternative Name:	Piccolo ( <a href="#">PIC Products</a> )
Background:	Piccolo, also referred to as Aczonin, is a large protein which consists of an N-terminal Zn <sup>2+</sup>

## Target Details

---

finger, several piccolo-bassoon homology domains and C-terminal PDZ and C2 domains (1, 2). In general it is found together with bassoon, a related huge multi-domain protein of the CAZ (cytoskeletal matrix assembled at active zones) (3). Piccolo is a scaffolding protein for proteins involved in endo- and exocytosis of synaptic vesicles. Piccolo has also been shown to interfere with clathrin mediated endocytosis by binding to the F-actin and dynamin binding protein Abp1 (2).

Gene ID: 26875

NCBI Accession: [NP\\_001104266](#)

UniProt: [Q9QYX7](#)

## Application Details

---

Application Notes:

- WB (1:1000)
- ICC/IF (1:100)
- optimal dilutions for assays should be determined by the user.

Comment: 1 µg/ml of ABIN863104 was sufficient for detection of Piccolo in 20 µg of rat brain tissue lysate by colorimetric immunoblot analysis using goat IgG:HRP as the secondary antibody.

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Concentration: 1 mg/mL


Buffer: PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: -20 °C

Storage Comment: -20°C



### Immunocytochemistry

**Image 1.** Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Piccolo Monoclonal Antibody, Clone 6H9-B6 (ABIN863104 and ABIN863105). Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-Piccolo Monoclonal Antibody (ABIN863104 and ABIN863105) at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:100 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain, DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60 min RT, 5 min RT. Localization: Cytoplasm, Endoplasmic Reticulum, Nucleus. Magnification: 60X. (A) DAPI (blue) nuclear stain. (B) Phalloidin Texas Red F-Actin stain. (C) Piccolo Antibody. (D) Composite.



### Western Blotting

**Image 2.** Western Blot analysis of Rat and mouse brain lysates showing detection of Piccolo protein using Mouse Anti-Piccolo Monoclonal Antibody, Clone 6H9-B6 (ABIN863104 and ABIN863105). Primary Antibody: Mouse Anti-Piccolo Monoclonal Antibody (ABIN863104 and ABIN863105) at 1:1000. (1) Mouse brain lysate, (2) Rat brain lysate.