antibodies -online.com





anti-KCNT1 antibody (AA 1168-1237)





Go to Product page

Overview

Quantity:	100 μg
Target:	KCNT1
Binding Specificity:	AA 1168-1237
Reactivity:	Rat
Host:	Mouse
Clonality:	Monoclonal
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC), Antibody Array (AA)

Product Details

Immunogen:	Fusion protein amino acids 1168-1237 of rat Slo2.2 (Slack)
Clone:	N3-26 (Formerly S3-26)
Isotype:	IgG1
Specificity:	Detects ~140 kDa. Weak human detection. Does not cross-react with KCNT2/Slo2.1/Slick.
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Protein G Purified

Target Details


Target:	KCNT1
Alternative Name:	Slo2.2 (KCNT1 Products)


Target Details

Storage Comment:

-20°C

l arget Details	
Background:	Slo2.2 is a novel member of the mammalian Slo potassium channel gene family. Slo2 channels may contribute to the resting potentials of cells that control their basal level of excitability (1). They also have sensors that couple channel activity to the intracellular concentrations of Na+ and Cl- (2).
Gene ID:	60444
NCBI Accession:	NP_068625
UniProt:	Q9Z258
Application Details	
Application Notes:	 WB (1:1000) IHC (1:1000) ICC/IF (1:100) optimal dilutions for assays should be determined by the user.
Comment:	1 μ g/ml of ABIN863116 was sufficient for detection of Slo2.2 in 10 μ g of rat brain lysate by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C





Immunocytochemistry

Image 1. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Slo2.2 Monoclonal Antibody, Clone N3/26 (ABIN863116). Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 4 % PFA for 15 min. Primary Antibody: Mouse Anti-Slo2.2 Monoclonal Antibody (ABIN863116) at 1:50 for overnight at 4 °C with slow rocking. Secondary Antibody: AlexaFluor 488 at 1:1000 for 1 hour at RT. Counterstain: Phalloidin-iFluor 647 (red) F-Actin stain, Hoechst (blue) nuclear stain at 1:800, 1.6 mM for 20 min at RT. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) Slo2.2 Antibody (D) Composite.

Western Blotting

Image 2. Slack Rat Brain Membrane Western Blotting.