

Datasheet for ABIN870667 anti-ULK3 antibody (C-Term)

1 Image



Go to Product page

_				
()	VA	rv	IPI	٨

Isotype:

Cross-Reactivity:

Purification:

Grade:

IgG

Mouse, Rat

Verified

Overview	
Quantity:	100 μg
Target:	ULK3
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This ULK3 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)
Product Details	
Purpose:	Ulk3 (mouse)
Immunogen:	Peptide with sequence C-KIRESHWEAESLDK, from the C Terminus of the protein sequence according to NP_001092906.1.
Sequence:	KIRESHWEAE SLDK

chromatography using the immunizing peptide.

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity

Target Details


Target:	ULK3
Alternative Name:	Ulk3 (ULK3 Products)
Background:	ULK3, unc-51-like kinase 3 (C. elegans), 1200015E14Rik, serine/threonine-protein kinase ULK3
Molecular Weight:	47kDa
Gene ID:	71742, 691171
NCBI Accession:	NP_082171

Application Details

Application Notes:	Immunohistochemistry: Paraffin embedded Human Kidney. Recommended concentration: 5 $\boldsymbol{\mu}$
	g/mL.
	Western Blot: Preliminary experiments gave an approx 47 kDa band in Rat Brain and Eye lysates
	after 1 µg/mL antibody staining. Please note that currently we cannot find an explanation in the
	literature for the band we observe given the calculated size of 5
	Peptide ELISA: antibody detection limit dilution 1:64000.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Minimize freezing and thawing.
Storage:	-20 °C
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable.



Immunohistochemistry

Image 1. ABIN870667 (5 μ g/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer Ph 6, AP-staining.