

[Go to Product page](#)

Datasheet for ABIN872355
anti-ZMIZ1 antibody (AA 51-150)

Overview

| | |
|----------------------|---|
| Quantity: | 100 µL |
| Target: | ZMIZ1 |
| Binding Specificity: | AA 51-150 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This ZMIZ1 antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA, Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |

Product Details

| | |
|-----------------------|--|
| Immunogen: | KLH conjugated synthetic peptide derived from human Retinoic acid induced 17 |
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse, Rat |
| Predicted Reactivity: | Dog,Cow,Sheep,Pig,Horse,Chicken |
| Purification: | Purified by Protein A. |

Target Details

| | |
|---------|-------|
| Target: | ZMIZ1 |
|---------|-------|

Target Details

| | |
|-------------------|--|
| Alternative Name: | Retinoic acid induced 17 (ZMIZ1 Products) |
| Background: | <p>Synonyms: Zinc finger MIZ domain-containing protein 1, MIZ, PIAS like protein hZimp10, PIAS like protein on chromosome 10, PIAS like protein Zimp10, PIAS-like protein Zimp10, RAI 17, RAI17, RAI-17, Retinoic acid induced 17, Retinoic acid induced protein 17, Retinoic acid-induced protein 17, TRAFIP10, Zimp10, Zinc finger containing Miz1 PIAS like protein on chromosome 10, Zinc finger MIZ domain-containing protein 1, Zinc finger MIZ-type containing 1, Zmiz1, ZMIZ1_HUMAN, hZIMP10, KIAA1224.</p> <p>Background: Increases ligand-dependent transcriptional activity of AR and promotes AR sumoylation. The stimulation of AR activity is dependent upon sumoylation. Tissue specificity: Expressed most abundantly in ovary and, at lower levels, in prostate, spleen and testis. Weak expression, if any, in thymus, small intestine, colon and peripheral blood leukocytes.</p> |
| Gene ID: | 57178 |

Application Details

| | |
|--------------------|---|
| Application Notes: | WB 1:300-5000 ELISA 1:500-1000 FCM 1:20-100 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 |
| Restrictions: | For Research Use only |

Handling

| | |
|--------------------|--|
| Format: | Liquid |
| Concentration: | 1 µg/µL |
| Buffer: | 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |

Handling

| | |
|----------|-------------|
| Storage: | 4 °C,-20 °C |
|----------|-------------|

| | |
|------------------|---|
| Storage Comment: | Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. |
|------------------|---|

| | |
|--------------|-----------|
| Expiry Date: | 12 months |
|--------------|-----------|