antibodies -online.com







anti-UBA1 antibody (AA 961-1058)



Image



Overview

Quantity:	100 μL
Target:	UBA1
Binding Specificity:	AA 961-1058
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This UBA1 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

. reader became	
Immunogen:	KLH conjugated synthetic peptide derived from human UBE1
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.
Target Details	

rarget Details

Target: UBA1


Target Details

_	
Alternative Name:	E1 Ubiquitin Activating Enzyme (UBA1 Products)
Background:	Synonyms: A1S9, A1S9 protein, A1S9T and BN75 temperature sensitivity complementing,
	A1S9T, A1ST, GXP 1, GXP1, MGC4781, Protein A1S9, Uba1, UBA1_HUMAN, UBE 1 , UBE 1X,
	UBE1, UBE1X, Ubiquitin activating enzyme E1, Ubiquitin-activating enzyme E1, Ubiquitin-like
	modier-activating enzyme 1.
	Background: The protein encoded by this gene catalyzes the first step in ubiquitin conjugation
	to mark cellular proteins for degradation. This gene complements an X-linked mouse
	temperature-sensitive defect in DNA synthesis, and thus may function in DNA repair. It is part of
	a gene cluster on chromosome Xp11.23. Alternatively spliced transcript variants that encode
	the same protein have been described. [provided by RefSeq, Jul 2008].
Gene ID:	7317
Application Details	
Application Notes:	ELISA 1:500-1000

Restrictions:	For Research Use only
	IF(ICC) 1:50-200
	IF(IHC-F) 1:50-200
	IF(IHC-P) 1:50-200
	IHC-F 1:100-500
	IHC-P 1:200-400
Application Notes:	ELISA 1:500-1000

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Paraformaldehyde-fixed, paraffin embedded rat brain, Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min, Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes, Blocking buffer (normal goat serum) at 37°C for 30min, Antibody incubation with E1 Ubiquitin Activating Enzyme Polyclonal Antibody, Unconjugated at 1:400 overnight at 4°C, followed by a conjugated secondary for 20 minutes and DAB staining.