antibodies -online.com





anti-Importin 4 antibody (AA 231-330)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL
Target:	Importin 4 (IPO4)
Binding Specificity:	AA 231-330
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Importin 4 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Importin4
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	Importin 4 (IPO4)
Alternative Name:	Importin4 (IPO4 Products)

Target Details	
Background:	Synonyms: Imp4, Imp4b, Importin 4b, Importin-4, Importin-4b, Ipo4, IPO4_HUMAN, Ran binding
	protein 4, Ran-binding protein 4, RanBP4, IPO4_HUMAN.
	Background: Functions in nuclear protein import as nuclear transport receptor. Serves as
	receptor for nuclear localization signals (NLS) in cargo substrates. Is thought to mediate
	docking of the importin/substrate complex to the nuclear pore complex (NPC) through binding
	to nucleoporin and the complex is subsequently translocated through the pore by an energy
	requiring, Ran-dependent mechanism. At the nucleoplasmic side of the NPC, Ran binds to the
	importin, the importin/substrate complex dissociates and importin is re-exported from the
	nucleus to the cytoplasm where GTP hydrolysis releases Ran. The directionality of nuclear
	import is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound
	forms of Ran between the cytoplasm and nucleus (By similarity). Mediates the nuclear import
	of RPS3A. In vitro, mediates the nuclear import of human cytomegalovirus UL84 by recognizing
	a non-classical NLS.
Gene ID:	79711
Pathways:	Protein targeting to Nucleus
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months