



[Go to Product page](#)

Datasheet for ABIN872401
anti-Nidogen antibody (AA 1001-1100)

Overview

Quantity:	100 µL
Target:	Nidogen (NID)
Binding Specificity:	AA 1001-1100
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Nidogen antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Entactin
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Cow,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	Nidogen (NID)
---------	---------------

Target Details

Alternative Name: [Nidogen \(NID Products\)](#)

Background: Synonyms: Ent, Entactin 1, Entactin, NID 1, NID, NID-1, Nid1, NID1_HUMAN, Nidogen enactin, Nidogen 1, Nidogen, Nidogen-1, NID1_HUMAN.

Background: Basement membranes are the earliest extracellular matrices produced during embryogenesis. They are synthesized and incorporated into the supramolecular architecture of several components, including laminins, Collagen IV, Nidogen and proteoglycans.

Nidogen/Entactin, a sulfated glycoprotein, acts as a link between the extracellular matrix molecules Laminin 1 and Collagen Type IV, and thereby participates in the assembly of basement membranes. Nidogen is a highly conserved member of the Nidogen family, which also includes Nidogen-2. Nidogen-2 has a high level of N- and O-glycosylation, and it interacts with Collagens Type I and IV and Perlecan at a comparable level to Nidogen. Nidogen is synthesized and secreted in primary and established mesenchymal peritubular cells and myoepithelial cells, and it affects adhesion of peritubular cells in an autocrine manner. Nidogen is expressed during embryonic and fetal development exclusively in fully developed basement membranes of the ectoderm and is not expressed in the developing endodermal basement membrane or in membranes disrupted during mesoderm formation. Nidogen also cooperates with Laminin 1 to regulate b-casein expression.

Gene ID: 4811

UniProt: [P14543](#)

Application Details

Application Notes: ELISA 1:500-1000
IHC-P 1:200-400
IHC-F 1:100-500
IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months