# antibodies - online.com







# anti-NPC1 antibody (AA 1181-1278)



#### Overview

Quantity:	100 μL
Target:	NPC1
Binding Specificity:	AA 1181-1278
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NPC1 antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human NPC1/Niemann Pick C1
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Pig,Chicken,Guinea Pig
Purification:	Purified by Protein A.

#### **Target Details**

Target: NPC1
--------------

## **Target Details**

Alternative Name:	NPC1 (NPC1 Products)
Background:	Synonyms: Niemann Pick C1, Niemann Pick C1 protein precursor, Niemann Pick disease, type
	C1, Niemann-Pick C1 protein, NPC, NPC1, NPC1_HUMAN.
	Background: This gene encodes a large protein that resides in the limiting membrane of
	endosomes and lysosomes and mediates intracellular cholesterol trafficking via binding of
	cholesterol to its N-terminal domain. It is predicted to have a cytoplasmic C-terminus, 13
	transmembrane domains, and 3 large loops in the lumen of the endosome - the last loop being
	at the N-terminus. This protein transports low-density lipoproteins to late endosomal/lysosoma
	compartments where they are hydrolized and released as free cholesterol. Defects in this gene
	cause Niemann-Pick type C disease, a rare autosomal recessive neurodegenerative disorder
	characterized by over accumulation of cholesterol and glycosphingolipids in late
	endosomal/lysosomal compartments.[provided by RefSeq, Aug 2009].
Gene ID:	4864
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

## Handling

Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months