antibodies -online.com





Datasheet for ABIN872436

anti-CEND1 antibody (AA 4-100)



Overview

Quantity:	100 μL
Target:	CEND1
Binding Specificity:	AA 4-100
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CEND1 antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CEND1
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Horse
Purification:	Purified by Protein A.
T 10.13	

Target Details

	Target:	CEND1		
--	---------	-------	--	--

Target Details

Alternative Name:	CEND1 (CEND1 Products)
Background:	Synonyms: BM88, BM88 antigen, Cell cycle exit and neuronal dferentiation 1, Cell cycle exit and
	neuronal dferentiation protein 1, FLJ90066, MGC34326, CEND_HUMAN
	Background: BM88 is a 149 amino acid protein that belongs to the CEND1 familly. Involved in
	neuroblastoma cell differentiation, BM88 is a single-pass type IV membrane protein that is
	neuron specific. It is suggested that BM88 forms a dimer of two identical polypeptides linked by
	disulfide bridges. BM88 has a central proline-rich region containing four PxxP motifs, which
	typically bind SRC homology-3 (SH3) domains, as well as a putative C-terminal transmembrane
	region, and several potential sites for N-glycosylation, myristoylation and phosphorylation. It is
	also suggested that a novel signaling mechanism exists by which BM88 interferes with calcium
	release from inositol 1,4,5-trisphosphate-sensitive stores and exerts anti-proliferative and anti-
	apoptotic functions. BM88 is an important molecular target for HDAC inhibition, and
	transcription of BM88 is induced by trichostatin-A.
Gene ID:	51286
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months