antibodies -online.com





anti-Nerve Growth Factor Receptor (TNFRSF16) Associated Protein 1 (NGFRAP1) (AA 34-111) antibody



Go to Product page

()	۱ ۱	\cap	r	/1	\cap	۱ ۸	1
0	'V	ㄷ	I١	νı	\Box	V	۷

Quantity:	100 μL
Target:	Nerve Growth Factor Receptor (TNFRSF16) Associated Protein 1 (NGFRAP1)
Binding Specificity:	AA 34-111
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	Un-conjugated
Application:	ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffinembedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NADE/NGFRAP1
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	Nerve Growth Factor Receptor (TNFRSF16) Associated Protein 1 (NGFRAP1)

Target Details

Alternative Name:	NGFRAP1 (NGFRAP1 Products)	
Background:	Synonyms: BEX3, BEX3_HUMAN, Brain expressed X linked 3 mouse homolog, Brain-expressed	
	X-linked protein 3, Nerve growth factor receptor TNFRSF16 associated protein 1, Nerve growth	
	factor receptor associated protein 1, Nerve growth factor receptor-associated protein 1,	
	NGFRAP1, Ovarian granulosa cell 13.0 kDa protein HGR74, Ovarian granulosa cell protein,	
	p75NTR associated cell death executor, p75NTR-associated cell death executor, Protein BEX3	
	Background: May be a signaling adapter molecule involved in p75NTR-mediated apoptosis	
	induced by NGF. Plays a role in zinc-triggered neuronal death (By similarity). May play an	
	important role in the pathogenesis of neurogenetic diseases. Tissue specificity: Found in	
	ovarian granulosa cells, testis, prostate and seminal vesicle tissue. High levels also detected in liver.	
Gene ID:	27018	
Pathways:	Neurotrophin Signaling Pathway, Positive Regulation of Endopeptidase Activity	
Application Details		
Application Notes:	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should b	
	handled by trained staff only.	
Storage:	4 °C,-20 °C	

Handling

Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months