# antibodies -online.com







# anti-NME5 antibody (AA 81-150)



#### Overview

Quantity:	100 μL	
Target:	NME5	
Binding Specificity:	AA 81-150	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This NME5 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human NME5/IPIA beta
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Horse,Rabbit
Purification:	Purified by Protein A.

## **Target Details**

Target:	NME5
Alternative Name:	NME5 (NME5 Products)

## **Target Details**

Storage:

Storage Comment:

rarget Details		
Background:	Synonyms: Inhibitor of p53 induced apoptosis beta, IPIA beta, NDK-H 5, NDP kinase homolog 5, NM23 H5, Nm23-M5, NM23H5, Non-metastatic cells 5, protein expressed in nucleoside-diphosphate kinase, Nonmetastatic cells 5, protein expressed in, Nonmetastatic protein 23, homolog 5, Nucleoside diphosphate kinase homolog 5, RSPH23, Testis specic nm23 homolog, NDK5_HUMAN.  Background: NME5 does not seem to have NDK kinase activity but confers protection from cell death by Bax and alters the cellular levels of several antioxidant enzymes including Gpx5. It may play a role in spermiogenesis by increasing the ability of late-stage spermatids to eliminate reactive oxygen species. NME5 is specifically expressed in testis germinal cells.	
Gene ID:	8382	
Pathways:	Nucleotide Phosphorylation, Ribonucleoside Biosynthetic Process, Negative Regulation of intrinsic apoptotic Signaling	
Application Details		
Application Notes:	WB 1:300-5000  ELISA 1:500-1000  IHC-P 1:200-400  IHC-F 1:100-500  IF(IHC-P) 1:50-200  IF(IHC-F) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

4 °C,-20 °C

$\vdash$	land	lına
	iaria	шц

Expiry Date:

12 months