antibodies -online.com







anti-CCDC43 antibody (AA 151-224)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL
Target:	CCDC43
Binding Specificity:	AA 151-224
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CCDC43 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CCDC43
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Pig,Horse
Purification:	Purified by Protein A.
Target Details	

Larget Details

Target:	CCDC43
Alternative Name:	CCDC43 (CCDC43 Products)

Target Details

Storage Comment:

rarget Details		
Background:	Synonyms: CCDC 43, Coiled coil domain containing 43, coiled coil domain containing protein	
	43, FLJ31795, CCD43_HUMAN.	
	Background: The coiled-coil domain is a structural motif found in proteins that are involved in a	
	diverse array of biological functions such as the regulation of gene expression, cell division,	
	membrane fusion and drug extrusion and delivery. CCDC43 (coiled-coil domain containing 43)	
	is a 224 amino acid protein encoded by a gene that maps to human chromosome 17q21.31.	
	Encoding over 1,200 genes, chromosome 17 comprises over 2.5 % of the human genome. Two	
	key tumor suppressor genes are associated with chromosome 17, namely, p53 and BRCA1.	
	Tumor suppressor p53 is necessary for maintenance of cellular genetic integrity by moderating	
	cell fate through DNA repair versus cell death. Malfunction or loss of p53 expression is	
	associated with malignant cell growth and Li-Fraumeni syndrome.	
Gene ID:	124808	
Application Details		
Application Notes:	WB 1:300-5000	
	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
-		

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

		1.	
\vdash	land	lın	
	ıaııu	1111	U

Expiry Date:

12 months