

Datasheet for ABIN872655 anti-DNASE1 antibody (AA 101-200)



Overview

Quantity:	100 μL
Target:	DNASE1
Binding Specificity:	AA 101-200
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DNASE1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DNase I
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:

Target Details

Alternative Name:	DNase 1 (DNASE1 Products)
Background:	Synonyms: RNASE1, Dornase alfa, Deoxyribonuclease 1, Deoxyribonuclease I,
	Deoxyribonuclease1, Deoxyribonucleasel, DNASE 1, DNase I lysosomal, DNASE1, DNasel, DNL
	1, DNL1, DRNI, Human urine deoxyribonuclease I, DNAS1_HUMAN.
	Background: Deoxyribonuclease I gene is approximately 3.2 kb long with 9 exons separated by
	8 introns. In the form of a bovine pancreatic enzyme preparation, it occupies an important place
	in the history of protein chemistry and enzymology: it was the first enzyme to be recognized as
	specific for DNA, it was the first DNase to be crystallized, and it was the first DNase for which a
	specific protein inhibitor was characterized. DNase I is a Ca2+ and Mg2+ dependant
	endonuclease. DNase I is synthesized in the pancreas and stored in zymogen granules. It has
	been used to reduce the viscosity of cystic fibrosis sputum. A DNase I-like enzyme appears to
	catalyze the degradation of chromatin to oligo- and mononucleosomes during apoptosis. A
	recent study has demonstrated an endonuclease with activity and antigenicity indistinguishable
	from DNase I in thymocytes, cells susceptible to apoptosis. DNase I is an endonuclease that
	hydrolyzes double-stranded or single stranded DNA preferentially at sites adjacent to pyrimidine
	nucleotides. The product of hydrolysis is a complex mixture of 5'-phosphate mononucleotides
	and oligonucleotides. In the presence of Mg ion, DNase I attacks each strand of DNA
	independently and the cleavage sites are random.
Gene ID:	1773
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
	Liquid
Format:	Liquid

Handling

Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months