# antibodies -online.com







## anti-MNAT1 antibody (AA 51-150)



( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μL
Target:	MNAT1
Binding Specificity:	AA 51-150
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MNAT1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human MNAT1
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

### Target Details

Target: MNAT1

### Target Details

Alternative Name:	MNAT1 (MNAT1 Products)	
Background:	Synonyms: MAT1, TFB3, CAP35, RNF66, CDK-activating kinase assembly factor MAT1,	
	CDK7/cyclin-H assembly factor, Cyclin-G1-interacting protein, Menage a trois, RING finger	
	protein 66, RING finger protein MAT1, p35, p36, MNAT1	
	Background: Stabilizes the cyclin H-CDK7 complex to form a functional CDK-activating kinase	
	(CAK) enzymatic complex. CAK activates the cyclin-associated kinases CDK1, CDK2, CDK4 and	
	CDK6 by threonine phosphorylation. CAK complexed to the core-TFIIH basal transcription	
	factor activates RNA polymerase II by serine phosphorylation of the repetitive C-terminal	
	domain (CTD) of its large subunit (POLR2A), allowing its escape from the promoter and	
	elongation of the transcripts. Involved in cell cycle control and in RNA transcription by RNA	
	polymerase II.	
Gene ID:	4331	
UniProt:	P51948	
Pathways:	Cell Division Cycle, Mitotic G1-G1/S Phases, M Phase	
Application Details		
Application Notes:	WB 1:300-5000	
	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	

### Handling

Storage:	4 °C,-20 °C
Storage Comment: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	
Expiry Date:	12 months