antibodies - online.com







anti-CORO1C antibody (AA 381-474)



Overview

Quantity:	100 μL
Target:	CORO1C
Binding Specificity:	AA 381-474
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CORO1C antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Troduct Details	
Immunogen:	KLH conjugated synthetic peptide derived from human Coronin 3
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.
Target Details	

rarget Details

Target: CORO1C

Target Details

Storage Comment:

Expiry Date:

rarget Details	
Alternative Name:	Coronin 3 (CORO1C Products)
Background:	Synonyms: CORO 1C, CORO1C, Coronin 1c, Coronin actin binding protein 1C, Coronin1c,
	Coronin3, CRNN 4, CRNN4, HCRNN 4, HCRNN4, COR1C_HUMAN.
	Background: Summary: This gene encodes a member of the WD repeat protein family. WD
	repeats are minimally conserved regions of approximately 40 amino acids typically bracketed
	by gly-his and trp-asp (GH-WD), which may facilitate formation of heterotrimeric or multiprotei
	complexes. Members of this family are involved in a variety of cellular processes, including cel
	cycle progression, signal transduction, apoptosis, and gene regulation. [provided by RefSeq, Ju
	2008].
Gene ID:	23603
Application Details	
Application Notes:	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	4 °C,-20 °C

12 months

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.