

Datasheet for ABIN872949  
**anti-PPM1F antibody (AA 325-425)**



[Go to Product page](#)

## Overview

|                      |  |
|----------------------|--|
| Quantity:            | 100 µL   |
| Target:              | PPM1F  |
| Binding Specificity: | AA 325-425   |
| Reactivity:          | Human  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This PPM1F antibody is un-conjugated   |
| Application:         | Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |

## Product Details

|                       |   |
|-----------------------|---|
| Immunogen:            | KLH conjugated synthetic peptide derived from human PPM1F |
| Isotype:              | IgG   |
| Predicted Reactivity: | Human,Mouse,Rat,Dog,Cow,Pig,Horse                         |
| Purification:         | Purified by Protein A.                                    |

## Target Details

|                   |  |
|-------------------|--|
| Target:           | PPM1F                                    |
| Alternative Name: | PPM1F ( <a href="#">PPM1F Products</a> ) |

## Target Details

|             |   |
|-------------|---|
| Background: | <p>Synonyms: Ca<sup>2+</sup>/calmodulin-dependent protein kinase phosphatase, CaM-kinase phosphatase, CaMKPase, hFEM-2, KIAA0015, Partner of PIX 2, POPX2, Protein phosphatase 1F, PPM1F_HUMAN.</p> <p>Background: The protein encoded by this gene is a member of the PP2C family of Ser/Thr protein phosphatases. PP2C family members are known to be negative regulators of cell stress response pathways. This phosphatase can interact with Rho guanine nucleotide exchange factors (PIX), and thus block the effects of p21-activated kinase 1 (PAK), a protein kinase mediating biological effects downstream of Rho GTPases. Calcium/calmodulin-dependent protein kinase II gamma (CAMK2G/CAMK-II) is found to be one of the substrates of this phosphatase. The overexpression of this phosphatase or CAMK2G has been shown to mediate caspase-dependent apoptosis. An alternatively spliced transcript variant has been identified, but its full-length nature has not been determined. [provided by RefSeq, Jul 2008].</p> |
| Gene ID:    | 9647  |
| Pathways:   | <a href="#">Positive Regulation of Endopeptidase Activity</a>   |

## Application Details

|                    |  |
|--------------------|--|
| Application Notes: | <p>WB 1:300-5000</p> <p>ELISA 1:500-1000</p> <p>IHC-P 1:200-400</p> <p>IHC-F 1:100-500</p> <p>IF(IHC-P) 1:50-200</p> <p>IF(IHC-F) 1:50-200</p> <p>IF(ICC) 1:50-200</p> |
| Restrictions:      | For Research Use only  |

## Handling

|                    |  |
|--------------------|--|
| Format:            | Liquid   |
| Concentration:     | 1 µg/µL  |
| Buffer:            | 0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.  |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |

Handling

---

|                  |   |
|------------------|---|
| Storage:         | 4 °C,-20 °C   |
| Storage Comment: | Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. |
| Expiry Date:     | 12 months   |