# antibodies -online.com





## anti-Cyclin L1 antibody (AA 351-450)



( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μL
Target:	Cyclin L1 (CCNL1)
Binding Specificity:	AA 351-450
Reactivity:	Chimpanzee
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Cyclin L1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human Cyclin L/CCNL1
Isotype:	IgG
Cross-Reactivity:	Chimpanzee
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Chimpanzee
Purification:	Purified by Protein A.

#### **Target Details**

Target: Cyclin L1 (CCNL1)
---------------------------

### Target Details

Alternative Name:	Cyclin L (CCNL1 Products)
Background:	Synonyms: Ania 6a, BM 001, CCNL 1, CCNL1, CCNL1_HUMAN, Cyclin L ania 6a, Cyclin L, Cyclin
	L gamma, Cyclin L1, Cyclin-L, Cyclin-L1.
	Background: Transcriptional regulator which participates in regulating the pre-mRNA splicing
	process. Seems to be involved in the regulation of RNA polymerase II (pol II). Functions in
	association with cyclin-dependent kinases (CDKs) and has a role in the second step of splicing.
	May be a candidate proto-oncogene in head and neck squamous cell carcinomas (HNSCC).
	Inhibited by the CDK-specific inhibitor p21. Tissue specificity, Ubiquitous with higher level in
	thymus. Overexpression in primary tumors of head and neck squamous cell carcinomas
	(HNSCC).
Gene ID:	57018

## Application Details

Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

$\vdash$	land	linc
	iaria	11116

Expiry Date:

12 months