antibodies - online.com







anti-DEAF1 antibody (AA 211-310)





)\/(

Overview	
Quantity:	100 μL
Target:	DEAF1
Binding Specificity:	AA 211-310
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DEAF1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human Deformed Epidermal Autoregulatory Factor 1
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Rat,Dog,Cow,Horse,Chicken
Purification:	Purified by Protein A.
Target Details	
Target:	DEAF1
Alternative Name:	Deformed Epidermal Autoregulatory Factor 1 (DEAF1 Products)
Alternative Name.	Deformed Epidermal Autoregulatory Factor 1 (DEALT Floducts)

Target Details

Bac	kar	ound:

Synonyms: DEAF1, Deformed epidermal autoregulatory factor 1 homolog, Nuclear DEA1 related transcriptional regulator, NUDR, SPN, Suppressin, Zinc finger MYND domain containing protein 5, ZMYND5, DEAF1_HUMAN.

Background: Suppressin, also known as DEAF1 is a transcription factor required for embryonic development. Suppressin contains one SAND domain and one C-terminal MYND-type zinc finger. It interacts with LMO4 and CLIM-2, suggesting that it plays a role mediating cell fate and embryonic pattern formation. Suppressin is expressed in a variety of tissues and localizes to the nucleus. Several isoforms exist due to alternative splicing and, depending on the isoform, Suppressin is secreted in some cell types. Secreted Suppressin can function to inhibit cell proliferation, arresting cells in the G0 or G1 phase. Mutations in the gene encoding Suppressin may result in a growth advantage leading to the development and progression of neoplasia. This suggest that Supressin is a potential target for cancer therapy.

Gene ID:

10522

Pathways:

Tube Formation

Application Details

Application Notes:

WB 1:300-5000

ELISA 1:500-1000


IF(IHC-P) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months



Western Blotting

Image 1. Mouse pancreas lysates, probed with Deformed Epidermal Autoregulatory Factor 1 Polyclonal Antibody, unconjugated (bs-6377R) at 1:2000 overnight at 4°C followed by a conjugated secondary antibody for 60 minutes at 37°C.