antibodies -online.com







anti-FRMD5 antibody (AA 211-312)



Image



\sim			
	$ \backslash / \cap$	r\/I	\square

Quantity:	100 μL
Target:	FRMD5
Binding Specificity:	AA 211-312
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FRMD5 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FRMD5	
Isotype:	IgG	
Cross-Reactivity:	Human	
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit	
Purification:	Purified by Protein A.	

Target Details

|--|--|--|


Target Details

Alternative Name:	Frmd5 (FRMD5 Products)	
Background:	Synonyms: FERM domain containing 5, FERM domain-containing protein 5, FLJ41022, FRMD5,	
	FRMD5_HUMAN, MGC14161.	
	Background: FERM domains are roughly 150 amino acids in length and are found in a number	
	of cytoskeletal-associated proteins such as Ezrin, Radixin, Moesin and 4.1 (erythrocyte	
	membrane protein band 4.1), where they provide a link between cytoskeletal signals and	
	membrane dynamics. FRMD5 (FERM domain-containing protein 5) is a 570 amino acid single-	
	pass membrane protein that contains one FERM domain and exists as two alternatively spliced	
	isoforms. The gene encoding FRMD5 maps to human chromosome 15, which houses over 700	
	genes and comprises nearly 3 % of the human genome. Angelman syndrome, Prader-Willi	
	syndrome, Tay-Sachs disease and Marfan syndrome are all associated with defects in	
	chromosome 15-localized genes.	
Gene ID:	84978	
Application Details		
Application Notes:	WB 1:300-5000	
	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	

Handling

Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months

Images



Western Blotting

Image 1. Lane 1: HepG2 lysates Lane 2: U937 lysates probed with FRMD5 Polyclonal Antibody, Unconjugated at 1:300 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at 1:10000 for 60 min at 37°C.